

# Creation and destruction of magnetic fields

Matthias Rempel

HAO/NCAR

July 30 2007

# Magnetic fields in the Universe

- Earth
  - Magnetic field present for  $\sim 3.5 \cdot 10^9$  years, much longer than Ohmic decay time ( $\sim 10^4$  years)
  - Strong variability on shorter time scales ( $10^3$  years)
- Mercury, Ganymede, (Io), Jupiter, Saturn, Uranus, Neptune have large scale fields
- Sun
  - Magnetic fields from smallest observable scales to size of sun
  - 11 year cycle of large scale field ([Movie](#))
  - Ohmic decay time  $\sim 10^9$  years (in absence of turbulence)
- Other stars
  - Stars with outer convection zone: similar to sun
  - Stars with outer radiation zone: most likely primordial fields
- Galaxies
  - Field structure coupled to observed matter distribution (e.g. spirals)
  - Only dynamo that is directly observable

# Scope of this lecture

- Processes of magnetic field generation and destruction in turbulent plasma flows
- Introduction to general concepts of dynamo theory (this is not a lecture about the solar dynamo!)
- Outline
  - MHD, induction equation
  - Some general remarks and definitions regarding dynamos
  - Large scale dynamos (mean field theory)
    - Kinematic theory
    - Characterization of possible dynamos
    - Non-kinematic effects
  - 3D simulations

# MHD equations

The full set of MHD equations combines the induction equation with the Navier-Stokes equations including the Lorentz-force:

$$\begin{aligned}\frac{\partial \varrho}{\partial t} &= -\nabla \cdot (\varrho \mathbf{v}) \\ \varrho \frac{\partial \mathbf{v}}{\partial t} &= -\varrho (\mathbf{v} \cdot \nabla) \mathbf{v} - \nabla p + \varrho \mathbf{g} + \frac{1}{\mu_0} (\nabla \times \mathbf{B}) \times \mathbf{B} + \nabla \cdot \boldsymbol{\tau} \\ \varrho \frac{\partial e}{\partial t} &= -\varrho (\mathbf{v} \cdot \nabla) e - p \nabla \cdot \mathbf{v} + \nabla \cdot (\kappa \nabla T) + Q_\nu + Q_\eta \\ \frac{\partial \mathbf{B}}{\partial t} &= \nabla \times (\mathbf{v} \times \mathbf{B} - \eta \nabla \times \mathbf{B})\end{aligned}$$

Assumptions:

- Validity of continuum approximation (enough particles to define averages)
- Non-relativistic motions, low frequencies
- Strong collisional coupling: validity of single fluid approximations, isotropic (scalar) gas pressure

# MHD equations

Viscous stress tensor  $\tau$

$$\begin{aligned}\Lambda_{ik} &= \frac{1}{2} \left( \frac{\partial v_i}{\partial x_k} + \frac{\partial v_k}{\partial x_i} \right) \\ \tau_{ik} &= 2\rho\nu \left( \Lambda_{ik} - \frac{1}{3} \delta_{ik} \nabla \cdot \mathbf{v} \right) \\ Q_\nu &= \tau_{ik} \Lambda_{ik} ,\end{aligned}$$

Ohmic dissipation  $Q_\eta$

$$Q_\eta = \frac{\eta}{\mu_0} (\nabla \times \mathbf{B})^2 .$$

Equation of state

$$p = \frac{\rho e}{\gamma - 1} .$$

$\nu$ ,  $\eta$  and  $\kappa$ : viscosity, magnetic diffusivity and thermal conductivity  
 $\mu_0$  denotes the permeability of vacuum

# Kinematic approach

- Solving the 3D MHD equations is not always feasible
- Semi-analytical approach preferred for understanding fundamental properties of dynamos
- Evaluate turbulent induction effects based on induction equation for a given velocity field
  - Velocity field assumed to be given as 'background' turbulence, Lorentz-force feedback neglected (sufficiently weak magnetic field)
  - What correlations of a turbulent velocity field are required for dynamo (large scale) action?
  - Theory of onset of dynamo action, but not for non-linear saturation
- More detailed discussion of induction equation

# Ohm's law

Equation of motion for drift velocity  $\mathbf{v}_d$  of electrons

$$m_e \left( \frac{\partial \mathbf{v}_d}{\partial t} + \frac{\mathbf{v}_d}{\tau_{ei}} \right) = -e(\mathbf{E} + \mathbf{v}_d \times \mathbf{B}) - \nabla p_e$$

$\tau_{ei}$ : collision time between electrons and ions

$-e$ : electron charge

$m_e$ : electron mass

$p_e$ : electron pressure

With the electric current:  $\mathbf{j} = -n e \mathbf{v}_d$  this gives the generalized Ohm's law:

$$\frac{\partial \mathbf{j}}{\partial t} + \frac{\mathbf{j}}{\tau_{ei}} = \frac{n_e e^2}{m_e} \mathbf{E} - \frac{e}{m_e} \mathbf{j} \times \mathbf{B} + \frac{n_e e}{m_e} \nabla p_e$$

Simplifications:

- $\tau_{ei} \omega_L \ll 1$ ,  $\omega_L = eB/m_e$ : Larmor frequency
- neglect  $\nabla p_e$
- low frequencies (no plasma oscillations)

Simplified Ohm's law

$$\mathbf{j} = \sigma \mathbf{E}$$

with the plasma conductivity

$$\sigma = \frac{\tau_{ei} n_e e^2}{m_e}$$

The Ohm's law we derived so far is only valid in the co-moving frame of the plasma. Under the assumption of non-relativistic motions this transforms in the laboratory frame to

$$\mathbf{j} = \sigma (\mathbf{E} + \mathbf{v} \times \mathbf{B})$$



# Induction equation\*

Using Ampere's law  $\nabla \times \mathbf{B} = \mu_0 \mathbf{j}$  yields for the electric field in the laboratory frame

$$\mathbf{E} = -\mathbf{v} \times \mathbf{B} + \frac{1}{\mu_0 \sigma} \nabla \times \mathbf{B}$$

leading to the induction equation

$$\frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \mathbf{E} = \nabla \times (\mathbf{v} \times \mathbf{B} - \eta \nabla \times \mathbf{B})$$

with the magnetic diffusivity

$$\eta = \frac{1}{\mu_0 \sigma} .$$

# Advection, diffusion, magnetic Reynolds number

$L$ : typical length scale  $U$ : typical velocity scale  $L/U$ : time unit

$$\frac{\partial \mathbf{B}}{\partial t} = \nabla \times \left( \mathbf{v} \times \mathbf{B} - \frac{1}{R_m} \nabla \times \mathbf{B} \right)$$

with the magnetic Reynolds number

$$R_m = \frac{U L}{\eta} .$$

$R_m \ll 1$ : diffusion dominated regime

$$\frac{\partial \mathbf{B}}{\partial t} = \eta \Delta \mathbf{B} .$$

Only decaying solutions with decay (diffusion) time scale

$$\tau_d \sim \frac{L^2}{\eta}$$

# Advection, diffusion, magnetic Reynolds number

$R_m \gg 1$  advection dominated regime (ideal MHD)

$$\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{v} \times \mathbf{B})$$

Equivalent expression

$$\frac{\partial \mathbf{B}}{\partial t} = -(\mathbf{v} \cdot \nabla) \mathbf{B} + (\mathbf{B} \cdot \nabla) \mathbf{v} - \mathbf{B} \nabla \cdot \mathbf{v}$$

- advection of magnetic field
- amplification by shear (stretching of field lines)
- amplification through compression



# Advection, diffusion, magnetic Reynolds number

Object	$\eta[\text{m}^2/\text{s}]$	$L[\text{m}]$	$U[\text{m/s}]$	$R_m$	$\tau_d$
earth (outer core)	2	$10^6$	$10^{-3}$	300	$10^4$ years
sun (plasma conductivity)	1	$10^8$	100	$10^{10}$	$10^9$ years
sun (turbulent conductivity)	$10^8$	$10^8$	100	100	3 years
liquid sodium lab experiment	0.1	1	10	100	10 s

# Alfven's theorem

Let  $\Phi$  be the magnetic flux through a surface  $F$  with the property that its boundary  $\partial F$  is moving with the fluid:

$$\Phi = \int_F \mathbf{B} \cdot d\mathbf{f} \longrightarrow \frac{d\Phi}{dt} = 0$$



- Flux is 'frozen' into the fluid
- Field lines 'move' with plasma

# Dynamos: Motivation

- For  $\mathbf{v} = 0$  magnetic field decays on timescale  $\tau_d \sim L^2/\eta$
- Earth and other planets:
  - Evidence for magnetic field on earth for  $3.5 \cdot 10^9$  years while  $\tau_d \sim 10^4$  years
  - Permanent rock magnetism not possible since  $T > T_{\text{Curie}}$  and field highly variable  $\longrightarrow$  field must be maintained by active process
- Sun and other stars:
  - Evidence for solar magnetic field for  $\sim 300\,000$  years ( $^{10}\text{Be}$ )
  - Most solar-like stars show magnetic activity independent of age
  - Indirect evidence for stellar magnetic fields over life time of stars
  - But  $\tau_d \sim 10^9$  years!
  - Primordial field could have survived in radiative interior of sun, but convection zone has much shorter diffusion time scale  $\sim 10$  years (turbulent diffusivity)

# Mathematical definition of dynamo

$S$  bounded volume with the surface  $\partial S$ ,  $\mathbf{B}$  maintained by currents contained within  $S$

$$\begin{aligned}\frac{\partial \mathbf{B}}{\partial t} &= \nabla \times (\mathbf{v} \times \mathbf{B} - \eta \nabla \times \mathbf{B}) && \text{in } S \\ \nabla \times \mathbf{B} &= 0 && \text{outside } S \\ [\mathbf{B}] &= 0 && \text{across } \partial S \\ \nabla \cdot \mathbf{B} &= 0\end{aligned}$$

$\mathbf{v} = 0$  outside  $S$ ,  $\mathbf{n} \cdot \mathbf{v} = 0$  on  $\partial S$  and

$$E_{\text{kin}} = \int_S \frac{1}{2} \rho \mathbf{v}^2 dV \leq E_{\text{max}} \quad \forall t$$

$\mathbf{v}$  is a dynamo if an initial condition  $\mathbf{B} = \mathbf{B}_0$  exists so that

$$E_{\text{mag}} = \int_{-\infty}^{\infty} \frac{1}{2\mu_0} \mathbf{B}^2 dV \geq E_{\text{min}} \quad \forall t$$

# Large scale/small scale dynamos

Decompose the magnetic field into large scale part and small scale part (energy carrying scale of turbulence)  $\mathbf{B} = \overline{\mathbf{B}} + \mathbf{B}'$ :


$$E_{\text{mag}} = \int \frac{1}{2\mu_0} \overline{\mathbf{B}}^2 dV + \int \frac{1}{2\mu_0} \overline{\mathbf{B}'^2} dV .$$

- Small scale dynamo:  $\overline{\mathbf{B}}^2 \ll \overline{\mathbf{B}'^2}$
- Large scale dynamo:  $\overline{\mathbf{B}}^2 \geq \overline{\mathbf{B}'^2}$

Almost all turbulent (chaotic) velocity fields are small scale dynamos for sufficiently large  $R_m$ , large scale dynamos require additional large scale symmetries (see second half of this lecture)



# Large scale/small scale dynamos



- Amplification through field line stretching
- Twist-fold required to repack field into original volume
- Magnetic diffusivity allows for change of topology

Influence of magnetic diffusivity on growth rate

- **Fast dynamo:** growth rate independent of  $R_m$   
(stretch-twist-fold mechanism)
- **Slow dynamo:** growth rate limited by resistivity  
(stretch-reconnect-repack)
- Fast dynamos relevant for most astrophysical objects since  $R_m \gg 1$
- Dynamos including (resistive) reconnection steps can be fast provided the reconnection is fast


# Differential rotation and meridional flow

Induction effects of axisymmetric flows on axisymmetric field:

$$\mathbf{B} = B\mathbf{e}_\Phi + \nabla \times (A\mathbf{e}_\Phi)$$

$$\mathbf{v} = v_r\mathbf{e}_r + v_\theta\mathbf{e}_\theta + \Omega r \sin\theta\mathbf{e}_\Phi$$

Differential rotation most dominant shear flow in stellar convection zones:



Meridional flow by-product of DR, observed as poleward surface flow in case of the sun

# Differential rotation and meridional flow

Spherical geometry:

$$\begin{aligned}\frac{\partial B}{\partial t} + \frac{1}{r} \left( \frac{\partial}{\partial r} (r v_r B) + \frac{\partial}{\partial \theta} (v_\theta B) \right) = \\ r \sin \theta \mathbf{B}_p \cdot \nabla \Omega + \eta \left( \Delta - \frac{1}{(r \sin \theta)^2} \right) B \\ \frac{\partial A}{\partial t} + \frac{1}{r \sin \theta} \mathbf{v}_p \cdot \nabla (r \sin \theta A) = \eta \left( \Delta - \frac{1}{(r \sin \theta)^2} \right) A\end{aligned}$$


- **Meridional flow:** Independent advection of poloidal and toroidal field
- **Differential rotation:** Source for toroidal field (if poloidal field not zero)
- **Diffusion:** Sink for poloidal and toroidal field
- No term capable of maintaining poloidal field against Ohmic decay!

# Differential rotation and meridional flow

- Weak poloidal seed field can lead to significant field amplification
  - No source term for poloidal field
  - Decay of poloidal field on resistive time scale
  - Ultimate decay of toroidal field
  - Not a dynamo!
  - What is needed?
- 
- Source for poloidal field

# Cowling's anti-dynamo theorem

A stationary axisymmetric magnetic field with currents limited to a finite volume in space cannot be maintained by a velocity field with finite amplitude.



Ohm's law of the form  $\mathbf{j} = \sigma \mathbf{E}$  only decaying solutions, focus here on  $\mathbf{j} = \sigma(\mathbf{v} \times \mathbf{B})$ .

On O-type neutral line  $\mathbf{B}_p$  is zero, but  $\mu_0 \mathbf{j}_t = \nabla \times \mathbf{B}_p$  has finite value, but cannot be maintained by  $(\mathbf{v} \times \mathbf{B})_t = (\mathbf{v}_p \times \mathbf{B}_p)$ .

# Large scale dynamo theory

Some history:

- 1919 Sir Joseph Larmor: Solar magnetic field maintained by motions of conducting fluid?
- 1937 Cowling's anti-dynamo theorem and many others
- 1955 Parker: decomposition of field in axisymmetric and non-axisymmetric parts, average over induction effects of non-axisymmetric field
- 1964 Braginskii, Steenbeck, Krause: Mathematical framework of mean field theory developed
- last 2 decades 3D dynamo simulations

# Reynolds rules

We need to define an averaging procedure to define the mean and the fluctuating field.

For any function  $f$  and  $g$  decomposed as  $f = \bar{f} + f'$  and  $g = \bar{g} + g'$  we require that the Reynolds rules apply

$$\begin{aligned}\overline{\bar{f}} &= \bar{f} \longrightarrow \overline{f'} = 0 \\ \overline{\bar{f} + \bar{g}} &= \bar{f} + \bar{g} \\ \overline{\bar{f} \bar{g}} &= \bar{f} \bar{g} \longrightarrow \overline{f' \bar{g}} = 0 \\ \overline{\partial f / \partial x_i} &= \partial \bar{f} / \partial x_i \\ \overline{\partial f / \partial t} &= \partial \bar{f} / \partial t .\end{aligned}$$

Examples:

- Longitudinal average (mean = axisymmetric component)
- Ensemble average (mean = average over several realizations of chaotic system)



# Meanfield induction equation

Average of induction equation:

$$\frac{\partial \bar{\mathbf{B}}}{\partial t} = \nabla \times \left( \overline{\mathbf{v}' \times \mathbf{B}'} + \bar{\mathbf{v}} \times \bar{\mathbf{B}} - \eta \nabla \times \bar{\mathbf{B}} \right)$$

New term resulting from small scale effects:

$$\bar{\mathcal{E}} = \overline{\mathbf{v}' \times \mathbf{B}'}$$

Fluctuating part of induction equation:

$$\left( \frac{\partial}{\partial t} - \eta \Delta \right) \mathbf{B}' - \nabla \times (\bar{\mathbf{v}} \times \mathbf{B}') = \nabla \times \left( \mathbf{v}' \times \bar{\mathbf{B}} + \mathbf{v}' \times \mathbf{B}' - \overline{\mathbf{v}' \times \mathbf{B}'} \right)$$

Kinematic approach:  $\mathbf{v}'$  assumed to be given

- Solve for  $\mathbf{B}'$ , compute  $\overline{\mathbf{v}' \times \mathbf{B}'}$  and solve for  $\bar{\mathbf{B}}$
- Term  $\mathbf{v}' \times \mathbf{B}' - \overline{\mathbf{v}' \times \mathbf{B}'}$  leading to higher order correlations (closure problem)

# Second order correlation approximation (SOCA)

Second order term can be neglected if

- $|\mathbf{B}'| \ll |\overline{\mathbf{B}}|$
- $|\mathbf{v}' \times \mathbf{B}' - \overline{\mathbf{v}' \times \mathbf{B}'}| \ll |\mathbf{v}' \times \overline{\mathbf{B}}|$
- $\nabla \times (\mathbf{v}' \times \mathbf{B}' - \overline{\mathbf{v}' \times \mathbf{B}'})$  correlates only weakly with  $\mathbf{v}'$

Sufficient condition:

- $R_m \ll 1$  or  $S = v\tau_c/l_c \ll 1 \longrightarrow |\mathbf{B}'| \ll |\overline{\mathbf{B}}|$
- Problem:  $R_m \gg 1$  and  $S \sim 1$  in stellar convection zones

In praxis it works better than it should!

# Second order correlation approximation (SOCA)

Neglecting higher order moments and assume  $\bar{\tau} \gg \tau_c$ :

$$\mathbf{B}' \approx \tau_c \nabla \times (\mathbf{v}' \times \bar{\mathbf{B}}) = -\tau_c (\mathbf{v}' \cdot \nabla) \bar{\mathbf{B}} + \tau_c [(\bar{\mathbf{B}} \cdot \nabla) \mathbf{v}' - \bar{\mathbf{B}} \nabla \cdot \mathbf{v}']$$

leads to the expression:

$$\bar{\mathcal{E}} = \alpha \bar{\mathbf{B}} + \gamma \times \bar{\mathbf{B}} - \beta \nabla \times \bar{\mathbf{B}} + \dots$$

with ( $\alpha$  and  $\beta$  are symmetric tensors):

$$\begin{aligned}\alpha_{ij} &= \frac{1}{2} \tau_c \left( \overline{\varepsilon_{ikl} v_{k'}' \frac{\partial v_{l'}'}{\partial x_j}} + \overline{\varepsilon_{jkl} v_{k'}' \frac{\partial v_{l'}'}{\partial x_i}} \right) \\ \gamma_i &= -\frac{1}{2} \tau_c \frac{\partial}{\partial x_k} \overline{v_i' v_k'} \\ \beta_{ij} &= \frac{1}{2} \tau_c \left( \overline{v'^2} \delta_{ij} - \overline{v_i' v_j'} \right)\end{aligned}$$

# Second order correlation approximation (SOCA)

Simplification for (quasi) isotropic, non-mirror symmetric, (weakly) inhomogeneous turbulence:

$$\overline{v_i' v_j'} \sim \delta_{ij}, \quad \alpha_{ij} = \alpha \delta_{ij}, \quad \beta_{ij} = \eta_t \delta_{ij}$$

Leads to:

$$\alpha = \frac{1}{3} \alpha_{ii} = -\frac{1}{3} \tau_c \overline{\mathbf{v}' \cdot (\nabla \times \mathbf{v}')} \sim \frac{\eta_t}{l_c} \sim v_{rms}$$

$$\eta_t = \frac{1}{3} \beta_{ii} = \frac{1}{3} \tau_c \overline{\mathbf{v}'^2} \sim l_c v_{rms}$$

$$\gamma = -\frac{1}{2} \nabla \eta_t$$

Induction equation for  $\overline{\mathbf{B}}$ :

$$\frac{\partial \overline{\mathbf{B}}}{\partial t} = \nabla \times [\alpha \overline{\mathbf{B}} + (\overline{\mathbf{v}} + \gamma) \times \overline{\mathbf{B}} - (\eta + \eta_t) \nabla \times \overline{\mathbf{B}}]$$

# Turbulent diffusivity - destruction of magnetic field

Turbulent diffusivity dominant dissipation process for large scale field in case of large  $R_m$ :

$$\eta_t = \frac{1}{3} \tau_c \overline{\mathbf{v}'^2} \sim L v_{\text{rms}} \sim R_m \eta \gg \eta$$

- Formally  $\eta_t$  comes from advection term (transport term, non-dissipative)
- Turbulent cascade transporting magnetic energy from the large scale  $L$  to the micro scale  $l_m$  (advection + reconnection)

$$\eta \mathbf{j}_m^2 \sim \eta_t \bar{\mathbf{j}}^2 \longrightarrow \frac{B_m}{l_m} \sim \sqrt{R_m} \frac{\bar{B}}{L}$$

**Important:** The large scale determines the energy dissipation rate,  $l$  adjusts to allow for the dissipation on the microscale.

Present for isotropic homogeneous turbulence

# Turbulent diamagnetism, turbulent pumping

Expulsion of flux from regions with larger turbulence intensity  
'diamagnetism'

$$\gamma = -\frac{1}{2}\nabla\eta_t$$

Downward directed at base of convection zone

Turbulent pumping (stratified convection):


- Upflows expand, downflows converge
- Stronger velocity and smaller filling factor of downflows
- Mean advection effect of up- and downflows does not cancel
- Downward transport found in numerical simulations

Requires inhomogeneity (stratification)

# Kinematic $\alpha$ -effect

$$\alpha = -\frac{1}{3}\tau_c \overline{\mathbf{v}' \cdot (\nabla \times \mathbf{v}')} \quad H_k = \overline{\mathbf{v}' \cdot (\nabla \times \mathbf{v}')} \quad \text{kinetic helicity}$$

Requires rotation + additional preferred direction (stratification)



# Fast or slow dynamo?

Turbulent induction effects require reconnection to operate; however, the expressions

$$\begin{aligned}\alpha_{ij} &= \frac{1}{2}\tau_c \left( \overline{\varepsilon_{ikl} v_k' \frac{\partial v_l'}{\partial x_j}} + \overline{\varepsilon_{jkl} v_k' \frac{\partial v_l'}{\partial x_i}} \right) \\ \gamma_i &= -\frac{1}{2}\tau_c \frac{\partial}{\partial x_k} \overline{v_i' v_k'} \\ \beta_{ij} &= \frac{1}{2}\tau_c \left( \overline{\mathbf{v}'^2} \delta_{ij} - \overline{v_i' v_j'} \right)\end{aligned}$$

are independent of  $\eta$  (in this approximation), indicating fast dynamo action.



# Validity of Mean field expansion

Second order correlation approximation:

- At best marginally justified
- Works better than it should

Most general form for mean field expansion:

$$\bar{\mathcal{E}}_i(\mathbf{x}, t) = \int_{-\infty}^{\infty} d^3x' \int_{-\infty}^t dt' \mathcal{K}_{ij}(\mathbf{x}, t, \mathbf{x}', t') \bar{\mathbf{B}}_j(\mathbf{x}', t') .$$

Sufficient **scale separation**

- $l_c \ll L$
- $\tau_c \ll \tau_L$

leads to:

$$\bar{\mathcal{E}} = \alpha \bar{\mathbf{B}} + \boldsymbol{\gamma} \times \bar{\mathbf{B}} - \beta \nabla \times \bar{\mathbf{B}} - \boldsymbol{\delta} \times \nabla \times \bar{\mathbf{B}} + \dots$$

In stellar convection zones scale separation also only marginally justified (continuous turbulence spectrum)!

**Large scale convection (M. Miesch, HAO)**

# Symmetry constraints

$\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$  depend on large scale symmetries of the system defining the symmetry properties of the turbulence (e.g. rotation and stratification). Additional to that the expansion

$$\overline{\mathcal{E}} = \alpha \overline{\mathbf{B}} + \gamma \times \overline{\mathbf{B}} - \beta \nabla \times \overline{\mathbf{B}} - \delta \times \nabla \times \overline{\mathbf{B}} + \dots$$

is a relation between polar and axial vectors:

- $\overline{\mathcal{E}}$ : polar vector, independent from handedness of coordinate system
- $\mathbf{B}$ : axial vector, involves handedness of coordinate system in definition (curl operator, cross product)

Handedness of coordinate system pure convention (contains no physics), consistency requires:

- $\alpha$ ,  $\delta$ : pseudo tensor
- $\beta$ ,  $\gamma$ : true tensors

# Symmetry constraints

## Turbulence with rotation and stratification

- true tensors:  $\delta_{ij}$ ,  $g_i$ ,  $g_i g_j$ ,  $\Omega_i \Omega_j$ ,  $\Omega_i \varepsilon_{ijk}$
- pseudo tensors:  $\varepsilon_{ijk}$ ,  $\Omega_i$ ,  $\Omega_i g_j$ ,  $g_i \varepsilon_{ijk}$

Symmetry constraints allow only certain combinations:

$$\begin{aligned}\alpha_{ij} &= \alpha_0 (\mathbf{g} \cdot \mathbf{\Omega}) \delta_{ij} + \alpha_1 (g_i \Omega_j + g_j \Omega_i) , & \gamma_i &= \gamma_0 g_i + \gamma_1 \varepsilon_{ijk} g_j \Omega_k \\ \beta_{ij} &= \beta_0 \delta_{ij} + \beta_1 g_i g_j + \beta_2 \Omega_i \Omega_j , & \delta_i &= \delta_0 \Omega_i\end{aligned}$$

The scalars  $\alpha_0 \dots \delta_0$  depend on quantities of the turbulence such as rms velocity and correlation times scale.

- isotropic turbulence: only  $\beta$
- + stratification:  $\beta + \gamma$
- + rotation:  $\beta + \delta$
- + stratification + rotation:  $\alpha$  can exist

# Generalized Ohm's law

What is needed to circumvent Cowling's theorem?


- Crucial for Cowling's theorem: Impossibility to drive a current parallel to magnetic field
- Cowling's theorem does not apply to mean field if a mean current can flow parallel to the mean field (since total field non-axisymmetric this is not a contradiction!)

$$\bar{\mathbf{j}} = \tilde{\sigma} (\bar{\mathbf{E}} + \bar{\mathbf{v}} \times \bar{\mathbf{B}} + \boldsymbol{\gamma} \times \bar{\mathbf{B}} + \alpha \bar{\mathbf{B}})$$

$\tilde{\sigma}$  contains contributions from  $\eta$ ,  $\beta$  and  $\delta$ .

Ways to circumvent Cowling:

- $\alpha$ -effect
- anisotropic conductivity (off diagonal elements +  $\delta$ -effect)



Induction of field parallel to current (producing helical field!)



$$\frac{\partial \bar{\mathbf{B}}}{\partial t} = \nabla \times (\alpha \bar{\mathbf{B}}) = \alpha \mu_0 \bar{\mathbf{j}}$$

Dynamo cycle:

$$\mathbf{B}_t \xrightarrow{\alpha} \mathbf{B}_p \xrightarrow{\alpha} \mathbf{B}_t$$

- Poloidal and toroidal field of similar strength
- In general stationary solutions

# $\alpha\Omega$ -, $\alpha^2\Omega$ -dynamo



Dynamo cycle:

$$\mathbf{B}_t \xrightarrow{\alpha} \mathbf{B}_p \xrightarrow{\Omega} \mathbf{B}_t$$

- Toroidal field much stronger than poloidal field
- In general traveling (along lines of constant  $\Omega$ ) and periodic solutions

$$\begin{aligned}\frac{\partial B}{\partial t} &+ \frac{1}{r} \left( \frac{\partial}{\partial r}(rv_r B) + \frac{\partial}{\partial \theta}(v_\theta B) \right) = r \sin \theta \mathbf{B}_p \cdot \nabla \Omega \\ &+ \eta \left( \Delta - \frac{1}{(r \sin \theta)^2} \right) B \\ \frac{\partial A}{\partial t} &+ \frac{1}{r \sin \theta} \mathbf{v}_p \cdot \nabla (r \sin \theta A) = \alpha B + \eta \left( \Delta - \frac{1}{(r \sin \theta)^2} \right) A\end{aligned}$$

- Dimensionless measure for strength of  $\Omega$ - and  $\alpha$ -effect


$$D_\Omega = \frac{R^2 \Delta \Omega}{\eta_t} \quad D_\alpha = \frac{R \alpha}{\eta_t}$$

- Dynamo excited if **dynamo number**

$$D = D_\Omega D_\alpha > D_{crit}$$

Movie:  $\alpha\Omega$ -dynamo

# $\alpha\Omega$ -dynamo with meridional flow



## Meridional flow:

- Poleward at top of convection zone
- Equatorward at bottom of convection zone

## Effect of advection:

- Equatorward propagation of activity
- Correct phase relation between poloidal and toroidal field
- Circulation time scale of flow sets dynamo period
- **Requirement:** Sufficiently low turbulent diffusivity

Movie: Flux-transport-dynamo (M. Dikpati, HAO)



$$\frac{\partial \bar{\mathbf{B}}}{\partial t} = \nabla \times [\delta \times (\nabla \times \bar{\mathbf{B}})] \sim \nabla \times (\Omega \times \bar{\mathbf{j}}) \sim \frac{\partial \bar{\mathbf{j}}}{\partial z}$$

- similar to  $\alpha$ -effect, but additional z-derivative of current
- couples poloidal and toroidal field
- $\delta^2$  dynamo is not possible:

$$\bar{\mathbf{j}} \cdot \bar{\mathcal{E}} = \bar{\mathbf{j}} \cdot (\delta \times \bar{\mathbf{j}}) = 0$$

- $\delta$ -effect is controversial (not all approximations give a non-zero effect)
- in most situations  $\alpha$  dominates

# Dynamos and magnetic helicity

Magnetic helicity (integral measure of field topology):

$$H_m = \int \mathbf{A} \cdot \mathbf{B} dV$$

has following conservation law (no helicity fluxes across boundaries):

$$\frac{d}{dt} \int \mathbf{A} \cdot \mathbf{B} dV = -2\mu_0 \eta \int \mathbf{j} \cdot \mathbf{B} dV$$

Decomposition into small and large scale part:

$$\begin{aligned} \frac{d}{dt} \int \overline{\mathbf{A}} \cdot \overline{\mathbf{B}} dV &= +2 \int \overline{\boldsymbol{\mathcal{E}}} \cdot \overline{\mathbf{B}} dV - 2\mu_0 \eta \int \overline{\mathbf{j}} \cdot \overline{\mathbf{B}} dV \\ \frac{d}{dt} \int \overline{\mathbf{A}'} \cdot \overline{\mathbf{B}'} dV &= -2 \int \overline{\boldsymbol{\mathcal{E}}} \cdot \overline{\mathbf{B}} dV - 2\mu_0 \eta \int \overline{\mathbf{j}'} \cdot \overline{\mathbf{B}'} dV \end{aligned}$$

# Dynamos and magnetic helicity

Dynamos have helical fields:

- $\alpha$ -effect induces magnetic helicity of same sign on large scale
- $\alpha$ -effect induces magnetic helicity of opposite sign on small scale

Saturation process (on scale  $\sim R_m \tau_c$ ):

$$\overline{\mathbf{j}' \cdot \mathbf{B}'} = -\bar{\mathbf{j}} \cdot \bar{\mathbf{B}} \longrightarrow \frac{|\bar{\mathbf{B}}|}{|\bar{\mathbf{B}}'|} \sim \sqrt{\frac{L}{l}}$$

$$\overline{\mathbf{j}' \cdot \mathbf{B}'} = -\frac{\alpha \bar{\mathbf{B}}^2}{\mu_0 \eta} + \frac{\eta_t}{\eta} \bar{\mathbf{j}} \cdot \bar{\mathbf{B}}$$

Time scales:

- Galaxy:  $\sim 10^{25}$  years ( $R_m \sim 10^{18}$ ,  $\tau_c \sim 10^7$  years)
- Sun:  $\sim 10^8$  years
- Earth:  $\sim 10^6$  years

# Non-kinematic effects

Proper way to treat them: 3D simulations

- Still very challenging
- Has been successful for geodynamo, but not for solar dynamo

Semi-analytical treatment of Lorentz-force feedback in mean field models:

- Macroscopic feedback: Change of the mean flow (differential rotation, meridional flow) through the mean Lorentz-force

$$\bar{\mathbf{f}} = \bar{\mathbf{j}} \times \bar{\mathbf{B}} + \overline{\mathbf{j}' \times \mathbf{B}'}$$

- Mean field model including mean field representation of full MHD equations:  
[Movie: Non-kinematic flux-transport dynamo](#)
- Microscopic feedback: Change of turbulent induction effects (e.g.  $\alpha$ -quenching)

Feedback of Lorentz force on small scale motions:

- Intensity of turbulent motions significantly reduced if  $\frac{1}{2\mu_0}B^2 > \frac{1}{2}\rho v_{rms}^2$ . Typical expression used

$$\alpha = \frac{\alpha_k}{1 + \frac{\overline{\mathbf{B}}^2}{B_{eq}^2}}$$

with the **equipartition field strength**  $B_{eq} = \sqrt{\mu_0 \rho} v_{rms}$

- Similar quenching also expected for turbulent diffusivity
- **Additional quenching of  $\alpha$  due to topological constraints possible (helicity conservation)**

**Controversial !**

# Microscopic feedback

Symmetry of momentum and induction equation  $\mathbf{v}' \leftrightarrow \mathbf{B}'$ :

$$\begin{aligned}\frac{d\mathbf{v}'}{dt} &= \frac{1}{\mu_0 \varrho} (\overline{\mathbf{B}} \cdot \nabla) \mathbf{B}' + \dots \\ \frac{d\mathbf{B}'}{dt} &= (\overline{\mathbf{B}} \cdot \nabla) \mathbf{v}' + \dots \\ \overline{\mathcal{E}} &= \overline{\mathbf{v}' \times \mathbf{B}'}\end{aligned}$$

Strongly motivates magnetic term for  $\alpha$ -effect (Pouquet et al. 1976):

$$\alpha = \frac{1}{3} \tau_c \left( \frac{1}{\varrho} \overline{\mathbf{j}' \cdot \mathbf{B}'} - \overline{\boldsymbol{\omega}' \cdot \mathbf{v}'} \right)$$

- Kinetic  $\alpha$ :  $\overline{\mathbf{B}} + \mathbf{v}' \longrightarrow \mathbf{B}' \longrightarrow \overline{\mathcal{E}}$
- Magnetic  $\alpha$ :  $\overline{\mathbf{B}} + \mathbf{B}' \longrightarrow \mathbf{v}' \longrightarrow \overline{\mathcal{E}}$

# Microscopic feedback

From helicity conservation one expects

$$\overline{\mathbf{j}' \cdot \mathbf{B}'} \sim -\alpha \overline{\mathbf{B}}^2$$

leading to algebraic quenching

$$\alpha = \frac{\alpha_k}{1 + g \frac{\overline{\mathbf{B}}^2}{B_{eq}^2}}$$

With the asymptotic expression (steady state)

$$\overline{\mathbf{j}' \cdot \mathbf{B}'} = -\frac{\alpha \overline{\mathbf{B}}^2}{\mu_0 \eta} + \frac{\eta_t}{\eta} \overline{\mathbf{j} \cdot \mathbf{B}}$$

we get

$$\alpha = \frac{\alpha_k + \frac{\eta_t^2}{\eta} \frac{\mu_0 \overline{\mathbf{j} \cdot \mathbf{B}}}{B_{eq}^2}}{1 + \frac{\eta_t}{\eta} \frac{\overline{\mathbf{B}}^2}{B_{eq}^2}}$$

# Microscopic feedback

Catastrophic  $\alpha$ -quenching ( $R_m \gg 1!$ ) in case of steady state and homogeneous  $\overline{\mathbf{B}}$ :

$$\alpha = \frac{\alpha_k}{1 + R_m \frac{\overline{\mathbf{B}}^2}{\overline{B}_{eq}^2}}$$

If  $\bar{\mathbf{j}} \cdot \overline{\mathbf{B}} \neq 0$  (dynamo generated field) and  $\eta_t$  unquenched:

$$\alpha \approx \eta_t \mu_0 \frac{\bar{\mathbf{j}} \cdot \overline{\mathbf{B}}}{\overline{\mathbf{B}}^2} \sim \frac{\eta_t}{L} \sim \frac{\eta_t}{l_c} \frac{l_c}{L} \sim \alpha_k \frac{l_c}{L}$$

- In general  $\alpha$ -quenching dynamic process: linked to time evolution of helicity
- Boundary conditions matter: Loss of small scale current helicity can alleviate catastrophic quenching
- Catastrophic  $\alpha$ -quenching turns large scale dynamo into slow dynamo



Stationary state reached on time scale  $R_m \tau_c$ :

- Galaxy:  $\sim 10^{25}$  years ( $R_m \sim 10^{18}$ ,  $\tau_c \sim 10^7$  years)
- Sun:  $\sim 10^8$  years
- Earth:  $\sim 10^6$  years
- Universe too young for galaxies to worry about stationary state!
- Sun, geodynamo had enough time too saturate
  - Sun: Possibility that helicity loss through photosphere alleviates quenching
  - Geodynamo:  $R_m \sim 300$  not that catastrophic?
- No observational evidence for catastrophic  $\alpha$ -quenching, but fundamental question for theory!

# 3D simulations

Why not just solving the full system to account for all non-linear effects?

- Most systems have  $R_e \gg R_m \gg 1$ , requiring high resolution
- Large scale dynamos evolve on time scales  $\tau_c \ll t \ll \tau_\eta$ , requiring long runs compared to convective turn over
- 3D simulations successful for geodynamo
  - $R_m \sim 300$ : all relevant magnetic scales resolvable
  - Incompressible system
- Solar dynamo: Ingredients can be simulated
  - Compressible system: density changes by  $10^6$  through convection zone
  - Boundary layer effects: Tachocline, difficult to simulate (strongly subadiabatic stratification, large time scales)
  - Magnetic structures down to 1000 km most likely important  
Evolve  $5000^3$  box over  $1000 \tau_c$ !
  - Small scale dynamos can be simulated (for  $P_m \sim 1$ )

# Summarizing remarks

## Destruction of magnetic field:

- Turbulent diffusivity: cascade of magnetic energy from large scale to dissipation scale (advection+reconnection)
- Enhances dissipation of large field by a factor  $R_m$

## Creation of magnetic field:

- Small scale dynamo (non-helical)
  - Amplification of field on and below scale of turbulence
  - Stretch-twist-fold-(reconnect)
  - Produces non-helical field and does not require helical motions
  - **Current research:** behavior for  $P_m \ll 1$
- Large scale dynamo (helical)
  - Amplification of field on scales larger than scale of turbulence
  - Produces helical field and does require helical motions
  - Requires rotation + additional symmetry direction  
(controversial  $\Omega \times J$  effect does not require helical motions)
  - **Current research:** catastrophic vs. non-catastrophic quenching